

EXAMPLES OF CLEAR LIQUID DIET



Einstein
Gastroenterology

—
Phone: 267.896.3745
—

COLONOSCOPY
PREP

Follow these directions ONLY

NULYTELY/GOLYTELY/COLYTE
Disregard directions on the preparation packaging

5 days BEFORE



Obtain PREP from your pharmacy



Review instructions regarding blood thinners (Plavix, Coumadin, Warfarin, etc). If you **DO NOT** understand these, call us as soon as possible.



STOP herbal, oil-based vitamins and iron supplements. **STOP** all medications that reduce diarrhea such as Imodium or Pepto-Bismol



Avoid salads or high fiber foods including **beans, nuts, seeds and popcorn** until after the procedure



Arrange for someone 18 or older to take you home after the procedure

1 day BEFORE



DRINK ONLY CLEAR LIQUIDS ALL DAY


- NO SOLID FOOD
- NO ALCOHOL
- NO RED OR PURPLE LIQUIDS



Examples of clear liquids include water, tea or coffee without cream, ginger ale, Sprite, apple juice, clear soup/broth, Gatorade, water-ice, Popsicles, Jell-O.



MIX PREP and place in fridge



DRINK WATER or clear liquids.
At least 1 cup (besides PREP) every 8 hours.




5 pm the night before
START drinking **HALF** of the PREP.
•Drink one cup (8 oz) every 15 minutes until half the gallon is consumed.

DAY of the procedure



6 hours prior to your procedure
Drink the second **HALF** of the PREP.
•This must be **FINISHED** at **LEAST 4** hours prior to your arrival time.



4 hours prior to your procedure
DO NOT DRINK OR EAT ANYTHING



You may take your medications with a sip of water.



Confirm that someone will be able to take you home after the procedure

